

Neuronal correlates of pitch in the Inferior Colliculus

Didier A. Depireux
David J. Klein
Jonathan Z. Simon
Shihab A. Shamma

Institute for Systems Research
University of Maryland
College Park, MD 20742-3311


*Supported in part by the Office of Naval Research, the National
Institute for Deafness and Communicative Disorders, and the National
Science Foundation.*

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2006		2. REPORT TYPE		3. DATES COVERED 00-00-2006 to 00-00-2006	
4. TITLE AND SUBTITLE Neuronal correlates of pitch in the Inferior Colliculus			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Maryland, Institute for Systems Research, College Park, MD, 20742-3311			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 18	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			


Methods

- ❑ Responses of single units in **Inferior Colliculus (IC)** and Primary Auditory Cortex (AI) in the **barbiturate-** or **ketamine-anesthetized ferret** were recorded with single tungsten electrodes. Data were collected from 13 ferrets, weighing 1.3 - 2.1 kg.
- ❑ **Surgery and Preparation:** The techniques involved are described in detail in Shamma et al. (1993). The ferrets were anesthetized with pentobarbital sodium and maintained in an areflexic state using a continuous IV infusion of pentobarbital or ketamine and xylazine, diluted with dextrose-electrolyte solution for metabolic stability. Data collection typically lasted 48-72 hours.
- ❑ **Recording Procedures:** Single-unit action potentials were recorded using glass-insulated tungsten microelectrodes with 5 to 6 M Ω impedance. The recorded signals were led through amplifiers and filters. Depending on the paradigm, a stimulus was presented every few seconds, and raster plots with time resolution of up to 0.1 ms were produced.
- ❑ IC was exposed by removal of (visual) cortex, and electrodes were lowered until ICC was reached, following standard criteria. Poorly defined best frequencies were very high at first, but went down very quickly as the electrode was lowered, corresponding to the ICX. When we reached the lowest Best Frequency (BF), corresponding to the top of the ICC, the responses changed qualitatively, and the BFs were better defined.

Auditory Pathway




The Inferior Colliculus



Why the IC?



- Midway up to Cortex
- Reports of IC maps and BMFs
- Observe good temporal responses



Theories of Pitch


Spectral

At minimum, there exists a resolved spectrum
 $200 + 400 + 600 + \dots + 2000 \text{ Hz}$




Temporal

No need for resolved spectrum
but **must** exist temporal properties of the response




Spectral Resolution & Ripples





Ripple Amplitude ΔA




Ripple phase Φ




Ripple frequency Ω



Response Field




Spectral Tuning to Ripples




Tuning to ripples based solely on Best Ripple Frequency indicates that cells' response areas are too broad to resolve harmonics.

Spectral Resolution & Ripples II



The modulation of the response to stationary ripples as a function of ripple phase decreases sharply as the ripple frequency increases, unlike in cortex. *Modulation* indicates the ratio of the maximum to the minimum response to a ripple of a given ripple frequency.

AM Rate Transfer Functions





Langner and Schreiner, *e.g.*, find that rate BMFs exhibit bandpass characteristics.



Langner and Schreiner (1988)

BMFs for AM Transfer Functions

Single cell
transfer
functions





Population
statistics of
transfer
functions



AM Transfer Function Characteristics

Synchronization




We characterize the AM synchronization transfer function by its peak or Best Modulation Frequency (BMF), as and upper cut-off, i.e. the frequency at which the synchronization coefficient is 50% of the peak value.


We find that the majority of cells have a BMF around 100 Hz, but with a range of cut-off frequencies.

Temporal Response to Pure Tones

Spike Train Autocorrelation




Fourier Transform




Temporal Response to AMs

Spike Train Autocorrelation




Fourier Transform




Temporal Response to Click Trains



Spike Train Autocorrelation





Fourier Transform



Fast Temporal Response I




2151C/02g09.k2p




Fast Temporal Response II

Raster of responses
to a click train.
Note that clicks'
phases are random
from sweep to sweep





215IC0812.k2

Autocorrelation
function for the
first four
frequencies



Inharmonic Stimulus



References

- ❑ DeValois R. and DeValois K. (1988) *Spatial Vision*. New York: Oxford U. Press.
- ❑ Langner G. and Schreiner C.E. (1988) "Periodicity coding in the inferior colliculus of the cat. I. Neuronal Mechanisms," *J. Neurophysiol.* 60(6), pp. 1799-1822.
- ❑ Langner G. (1992) "Periodicity coding in the auditory system," *Hear. Res.* 60, pp. 115-142.
- ❑ Shamma S.A., Versnel H. and Kowalski N. (1995) "Ripple analysis in ferret primary auditory cortex I. Response characteristics of single units to sinusoidally rippled spectra," *Auditory Neuroscience* 1(3), pp. 233-254.
- ❑ Shamma S.A. and Versnel H. (1995) "Ripple analysis in ferret primary auditory cortex. II. Prediction of unit responses to arbitrary spectral profiles," *Auditory Neuroscience* 1(3), pp. 255-270.
- ❑ Versnel H., Kowalski N. and Shamma S.A. (1995) "Ripple analysis in ferret primary auditory cortex. III. Topographic distribution of ripple response parameters," *Auditory Neuroscience* 1(3), pp. 271-285.
- ❑ Schreiner C.E. and Calhoun B.M. (1995) "Spectral envelope coding in cat primary auditory cortex: properties of ripple transfer functions," *Auditory Neuroscience* 1(1), 23 pages.
- ❑ Kowalski N., Depireux D.A. and Shamma S.A. (1996) "Analysis of dynamic spectra in ferret primary auditory cortex: I. Response characteristics of single units to moving rippled spectra," *J. Neurophysiol.* 76(5), pp. 3503-3523.
- ❑ Kowalski N., Depireux D.A. and Shamma S.A. (1996) "Analysis of dynamic spectra in ferret primary auditory cortex: II. Prediction of unit responses to arbitrary dynamic spectra," *J. Neurophysiol.* 76(5), pp. 3524-3534.